

CONTRACTIONS AND DEFORMATIONS

WILL DONOVAN AND MICHAEL WEMYSS

ABSTRACT. Suppose that f is a projective birational morphism with at most one-dimensional fibres between d -dimensional varieties X and Y , satisfying $\mathbf{R}f_*\mathcal{O}_X = \mathcal{O}_Y$. Consider the locus L in Y over which f is not an isomorphism. Taking the scheme-theoretic fibre C over any closed point of L , we construct algebras A_{fib} and A_{con} which prorepresent the functors of commutative deformations of C , and noncommutative deformations of the reduced fibre, respectively. Our main theorem is that the algebras A_{con} recover L , and in general the commutative deformations of neither C nor the reduced fibre can do this. As the $d = 3$ special case, this proves the following contraction theorem: in a neighbourhood of the point, the morphism f contracts a curve without contracting a divisor if and only if the functor of noncommutative deformations of the reduced fibre is representable.

CONTENTS

1. Introduction	1
2. Naive and DG Deformations	4
3. Deformations of Reduced Fibres	8
4. The Contraction Theorem	11
5. Deformations of the Scheme-Theoretic Fibre	15
6. Examples	16
References	19

1. INTRODUCTION

Our setting is a contraction $f: X \rightarrow X_{\text{con}}$ with at most one-dimensional fibres between d -dimensional varieties, satisfying $\mathbf{R}f_*\mathcal{O}_X = \mathcal{O}_{X_{\text{con}}}$. Writing $L \subset X_{\text{con}}$ for the locus over which f is not an isomorphism, it is a fundamental problem to characterise L , locally around a closed point in the base. For this, it is natural to study the deformations of the curve(s) above the point, and the question which we answer in this paper is the following.

Question. Which deformation-theoretic framework detects the non-isomorphism locus L , Zariski locally around a closed point $p \in X_{\text{con}}$?

The answer turns out to lie in noncommutative deformations, without assumptions on the singularities of X , and in arbitrary dimension. This process associates a noncommutative algebra to each point, which should be viewed as an invariant of the contraction f . When $d = 3$ and the algebra is finite-dimensional, its dimension has a curve-counting interpretation [T], but in all cases the algebra structure gives information about the neighbourhood of the point. This extra information has applications in constructing derived autoequivalences [DW1, DW3, W, K], and also in the minimal model program, allowing us to control iteration of flops and count minimal models [W], and produce the first explicit examples of Type E flops [BW]. It is also conjectured that such algebras classify smooth 3-fold flops [DW1, HT], and furthermore it is expected that they control divisor-to-curve contractions.

2010 *Mathematics Subject Classification.* Primary 14D15; Secondary 14E30, 14F05, 16E45, 16S38. The first author was supported by World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan, and by EPSRC grant EP/G007632/1. The second author was supported by EPSRC grant EP/K021400/1.




FIGURE 1. Contractions of 3-folds: divisor to curve, and curve to point.

1.1. Summary of Results. For a closed point $p \in L$, consider the scheme-theoretic fibre $C = f^{-1}(p)$. Set-theoretically, it is well known that C is a union of \mathbb{P}^1 s, and we denote these by C_1, \dots, C_n . To specify a deformation problem requires us to provide test objects, and to say what object(s) are being deformed. The test objects for the noncommutative deformation functor are the category \mathbf{Art}_n of artinian augmented \mathbb{C}^n -algebras, and the objects being deformed are $\{\mathcal{O}_{C_1}(-1), \dots, \mathcal{O}_{C_n}(-1)\}$. Informally, we wish to control the deformations of the $\mathcal{O}_{C_i}(-1)$, and also the mutual extensions between them. Formally, as explained in §2, this is encoded via the Maurer–Cartan formulation as a functor

$$\mathcal{Def}^{\mathcal{J}} : \mathbf{Art}_n \rightarrow \mathbf{Sets}.$$

Given a closed point $p \in X_{\text{con}}$, it is well-known that the formal fibre over p is derived equivalent to a certain noncommutative ring A . By taking suitable factors, as in [DW1, §2] we obtain the contraction algebra A_{con} , referring the reader to 3.5 for full details. Our first main result is then the following. The special case where the locus L is a single point, $d = 3$, and $n = 1$, was previously shown in [DW1, 3.1].

Theorem 1.1 (=3.9). *For each closed point $p \in L$, the algebra A_{con} (depending on p) prorepresents the functor of noncommutative deformations $\mathcal{Def}^{\mathcal{J}}$ of the reduced fibre over p .*

It turns out that the geometry of the locus L is controlled by the support of A_{con} .

Theorem 1.2. *With the setup above, pick an affine open neighbourhood $\text{Spec } R$ in X_{con} , and consider $L_R := L \cap \text{Spec } R$.*

- (1) (=3.3, 4.7(1)) *There is an R -algebra Λ_{con} which satisfies $\text{Supp}_R \Lambda_{\text{con}} = L_R$.*
- (2) (=3.7) *For each closed point $p \in L_R$, the completion of Λ_{con} at p is morita equivalent to A_{con} .*

This theorem has two main consequences.

Corollary 1.3. *The dimension of L_R at p is $\dim_{\mathfrak{R}} \text{Supp}_{\mathfrak{R}} A_{\text{con}}$, where \mathfrak{R} denotes the completion of R at p .*

Theorem 1.4 (=4.8, contraction theorem). *When $d = 3$, there is a neighbourhood of p over which f does not contract a divisor if and only if $\dim_{\mathbb{C}} A_{\text{con}} < \infty$.*

The ‘only if’ direction is easy and is known from our previous work [DW1, 2.13]. The content is the ‘if’ direction, and this requires significantly more technology.

1.2. Comparing deformation theories. We next show that other natural deformation functors do not control the geometry of L . As above, consider the scheme-theoretic fibre $C = f^{-1}(p)$ and the reduced curves C_1, \dots, C_n therein. To this data, we associate three other deformation problems. Again the details are left to §2, but the following table

summarises all four functors, giving the test objects and the deformed object in each case. Here \mathbf{CArt}_n is the category of commutative artinian augmented \mathbb{C}^n -algebras.

Deformation problem	Functor	Test objects	Object(s) deformed
Classical scheme-theoretic	$cDef^{\mathcal{O}_C}$	\mathbf{CArt}_1	\mathcal{O}_C
Noncommutative scheme-theoretic	$Def^{\mathcal{O}_C}$	\mathbf{Art}_1	\mathcal{O}_C
Commutative multi-pointed	$cDef^{\mathcal{I}}$	\mathbf{CArt}_n	$\oplus_i \mathcal{O}_{C_i}(-1)$
Noncommutative multi-pointed	$Def^{\mathcal{I}}$	\mathbf{Art}_n	$\{\mathcal{O}_{C_i}(-1)\}_i$

The following result drops out of our general construction. It is quite surprising, since it says that noncommutative deformations of the scheme-theoretic fibre give nothing in addition to the classical ones.

Theorem 1.5 (=5.3). *The functors $cDef^{\mathcal{O}_C}$ and $Def^{\mathcal{O}_C}$ are prorepresented by the same object A_{fib} .*

In general, however, the prorepresenting objects for the functors $cDef^{\mathcal{O}_C}$, $cDef^{\mathcal{I}}$ and $Def^{\mathcal{I}}$ are different. We prove the following.

Proposition 1.6 (=6.3, 6.4). *Neither $cDef^{\mathcal{O}_C}$ nor $cDef^{\mathcal{I}}$ detect the dimension of the non-isomorphism locus L .*

This is clear when the fibre above p has more than one irreducible curve, but is much more surprising when there is only a single irreducible curve in the fibres. We produce in 6.4 a contraction, sketched below, with a one-dimensional non-isomorphism locus L in which the central point 0 is cD_4 , and all other points are cA_1 . The commutative deformations of all reduced fibres except the central one are infinite dimensional, whereas the noncommutative deformations are always infinite dimensional.




FIGURE 2. Commutative versus noncommutative deformations of C^{red} .

Thus, it follows that $Def^{\mathcal{I}}$ is the functor that controls the contractibility of curves. For the convenience of the reader, we summarise the above together with the main results of [DW3] in the following table. To discuss autoequivalences, we further assume that f is a flopping contraction, and X is \mathbb{Q} -factorial with only Gorenstein terminal singularities.

Deformation problem	Functor	Detects divisor?	Corresponds to autoequivalence?
Classical	$cDef^{\mathcal{O}_C}$	No	Yes
Commutative multi-pointed	$cDef^{\mathcal{I}}$	No	No
Noncommutative multi-pointed	$Def^{\mathcal{I}}$	Yes	Yes

Our method to prove the above theorems involves noncommutative deformation theory associated to DGAs. By passing through various derived equivalences and embeddings, we reduce the geometric deformation problem into an easier problem about simultaneously deforming a collection of simple modules on a complete local ring obtained by tilting.

1.3. Structure of the Paper. In §2.1, we recall the naive noncommutative deformation theory developed by Laudal [L02], and subsequently Eriksen [E07]. We then describe DG deformation theory in §2.2, noting that it is equivalent by work of Segal [S08] and Efimov–Lunts–Orlov [ELO], and establish tools for comparing DG deformations which will arise in our construction.

In §3 we give the geometric setup, before proving the prorepresentability results for noncommutative deformations of the reduced fibre. The key observation, building on §2, is that this deformation functor on X is isomorphic to a DG deformation functor associated to a specific locally free resolution on U , naturally obtained via tilting. In §4 we use these results to prove 1.1, 1.2, and the Contraction Theorem.

In §5 we prove the prorepresentability results for both commutative and noncommutative deformations of the scheme-theoretic fibre, and show that they are prorepresented by the same object. The fact that A_{con} and A_{fib} are obtained as factors of a common ring A then allows us to relate the different deformation functors, and we do this in §5.2. We conclude in §6 by giving examples which illustrate the necessity of noncommutative deformations in the above theorems.

1.4. Conventions. Throughout we work over the field of complex numbers \mathbb{C} . Unqualified uses of the word ‘module’ refer to left modules, and $\text{mod } A$ denotes the category of finitely generated left A -modules. For two \mathbb{C} -algebras A and B , an A - B bimodule is the same thing as an $A \otimes_{\mathbb{C}} B^{\text{op}}$ -module. We use the functorial convention for composing arrows, so $f \cdot g$ means f then g . In particular, naturally this makes $M \in \text{mod } A$ into an $\text{End}_R(M)$ -module. We remark that these conventions are opposite to those in [DW1, DW3], but we do this to match the conventions in [E07, ELO]. Similarly, for quivers, DG category morphisms, and matrix multiplication, we write ab for a then b . We reserve the notation $f \circ g$ for the composition g then f .

In an abelian category \mathcal{A} , given two objects $a, b \in \mathcal{A}$ where a is a summand of b , we write $[a]$ for the two-sided ideal of $\text{End}(b)$ consisting of all morphisms that factor through a summand of a finite direct sum of a ’s.

Acknowledgements. The authors would like to thank Jon Pridham, Ed Segal, Olaf Schnürer and Yukinobu Toda for helpful discussions relating to this work.

2. NAIVE AND DG DEFORMATIONS

This section is mainly a review of known material, and is used to set notation. Noncommutative deformations of modules were introduced by Laudal [L02], and we review these naive deformation functors in §2.1 below. In our geometric setting later, this naive definition is necessary in order to establish the prorepresenting object in 3.9.

However, it is cumbersome to compare two or more naive deformation functors, as is demonstrated in the proofs of [DW1, 2.6, 2.8, 2.19], and for this the setting of multi-pointed DG deformation functors is much better suited. We review this theory in §2.2, being a slight generalisation of the setting of [S08, ELO], before in §2.3 proving some general DG deformation functor results.

2.1. Naive Deformations of Modules. From the geometric motivation of the introduction, where we want to deform n reduced curves in a fibre simultaneously, the test objects for the naive deformation functors are objects of the category \mathbf{Art}_n , defined as follows.

Definition 2.1. *An n -pointed \mathbb{C} -algebra Γ is an associative \mathbb{C} -algebra, together with \mathbb{C} -algebra morphisms $p: \Gamma \rightarrow \mathbb{C}^n$ and $i: \mathbb{C}^n \rightarrow \Gamma$ such that $ip = \text{Id}$. A morphism of n -pointed*

\mathbb{C} -algebras $\psi: (\Gamma, p, i) \rightarrow (\Gamma', p', i')$ is an algebra homomorphism $\psi: \Gamma \rightarrow \Gamma'$ such that

$$\begin{array}{ccccc} & & \Gamma & & \\ & i \nearrow & \downarrow \psi & \searrow p & \\ \mathbb{C}^n & & & & \mathbb{C}^n \\ & i' \searrow & \downarrow \psi & \nearrow p' & \\ & & \Gamma' & & \end{array}$$

commutes. We denote the category of n -pointed \mathbb{C} -algebras by \mathbf{Alg}_n . We denote the full subcategory consisting of those objects that are commutative rings by \mathbf{CAlg}_n .

We write \mathbf{Art}_n for the full subcategory of \mathbf{Alg}_n consisting of objects (Γ, p, i) for which $\dim_{\mathbb{C}} \Gamma < \infty$ and the augmentation ideal $\mathfrak{n} := \text{Ker } p$ is nilpotent. We write \mathbf{CArt}_n for the full subcategory of \mathbf{Art}_n consisting of those objects that are commutative rings.

Remark 2.2. If Γ is an associative ring, then by [E03, 1.3], there exists p, i such that $(\Gamma, p, i) \in \mathbf{Art}_n$ if and only if Γ is an artinian \mathbb{C} -algebra with precisely n simple modules (up to isomorphism), each of them one-dimensional over \mathbb{C} .

Notation 2.3. If $(\Gamma, p, i) \in \mathbf{Art}_n$, the structure morphisms p and i allow us to lift the canonical idempotents $\{e_i, \dots, e_n\}$ of \mathbb{C}^n to Γ . We will write

$$\Gamma_{ij} := e_i \Gamma e_j,$$

and consider Γ as a matrix ring (Γ_{ij}) under standard matrix multiplication. Accordingly, a left Γ -module M may be described in terms of its vector of summands $M_i := e_i M$, and a right Γ -module N can be described by its summands $N_i := N e_i$.

Definition 2.4. Given a \mathbb{C} -algebra Λ , choose a family $\mathcal{S} = \{S_1, \dots, S_n\}$ of objects in $\text{Mod } \Lambda$. The deformation functor

$$\text{Def}_{\Lambda}^{\mathcal{S}}: \mathbf{Art}_n \rightarrow \mathbf{Sets}$$

is defined by sending

$$(\Gamma, \mathfrak{n}) \mapsto \left\{ (M, \underline{\delta}) \left| \begin{array}{l} M \in \text{Mod } \Lambda \otimes_{\mathbb{C}} \Gamma^{\text{op}} \\ M \cong (S_i \otimes_{\mathbb{C}} \Gamma_{ij}) \text{ as right } \Gamma\text{-modules} \\ \underline{\delta} = (\delta_i), \text{ where each } \delta_i: M \otimes_{\Gamma} (\Gamma/\mathfrak{n})e_i \xrightarrow{\sim} S_i \end{array} \right. \right\} / \sim$$

where

- (1) $(S_i \otimes_{\mathbb{C}} \Gamma_{ij})$ refers to the \mathbb{C} -vector space $(S_i \otimes_{\mathbb{C}} \Gamma_{ij}) := \bigoplus_{i,j=1}^n (S_i \otimes_{\mathbb{C}} \Gamma_{ij})$, with the natural right Γ -module structure coming from the multiplication in Γ .
- (2) $(M, \underline{\delta}) \sim (N, \underline{\delta}')$ iff there exists an isomorphism $\tau: M \rightarrow N$ of bimodules such that the following diagram commutes for all $i = 1, \dots, n$.

$$\begin{array}{ccc} M \otimes_{\Gamma} (\Gamma/\mathfrak{n})e_i & \xrightarrow{\tau \otimes 1} & N \otimes_{\Gamma} (\Gamma/\mathfrak{n})e_i \\ & \searrow \delta_i & \swarrow \delta'_i \\ & S_i & \end{array}$$

An important problem in deformation theory is determining when deformation functors are prorepresentable, and also effectively describing the prorepresenting object. We briefly recall these notions, mainly to fix notation.

For any $(\Gamma, p, i) \in \mathbf{Alg}_n$, setting $I(\Gamma) := \text{Ker } p$ we consider the $I(\Gamma)$ -adic completion $\widehat{\Gamma}$ of Γ , defined by

$$\widehat{\Gamma} := \varprojlim \Gamma/I(\Gamma)^n.$$

The canonical morphism $\psi_{\Gamma}: \Gamma \rightarrow \widehat{\Gamma}$ belongs to \mathbf{Alg}_n . We say that $\Gamma \in \mathbf{Alg}_n$ is *complete* if ψ_{Γ} is an isomorphism. The pro-category \mathbf{pArt}_n is then defined to be the full subcategory of \mathbf{Alg}_n consisting of those objects (Γ, p, i) for which Γ is $I(\Gamma)$ -adically complete, and $\Gamma/I(\Gamma)^r \in \mathbf{Art}_n$ for all $r \geq 1$. It is clear that $\mathbf{Art}_n \subseteq \mathbf{pArt}_n$.

For a deformation functor $F: \mathbf{Art}_n \rightarrow \mathbf{Sets}$, recall that

- (1) F is called *prorepresentable* if $F \cong \text{Hom}_{\mathbf{pArt}_n}(\Gamma, -)|_{\mathbf{Art}_n}$ for some $\Gamma \in \mathbf{pArt}_n$.
- (2) F is called *representable* if $F \cong \text{Hom}_{\mathbf{Art}_n}(\Gamma, -)$ for some $\Gamma \in \mathbf{Art}_n$.

It is clear that if F is prorepresented by $\Gamma \in \mathbf{pArt}_n$, then F is representable if and only if $\dim_{\mathbb{C}} \Gamma < \infty$.

It is well-known that the functor in 2.4 is prorepresentable if $\mathrm{Ext}_{\Lambda}^t(\bigoplus S_i, \bigoplus S_i)$ is finite dimensional for $t = 1, 2$ [L02]: we will not use this fact below, however, instead preferring to establish the prorepresenting object in a much more direct way.

2.2. DG Deformations. With our conventions as in the introduction, recall that a DG category is a graded category \mathbf{A} whose morphism spaces are endowed with a differential d , i.e. homogeneous maps of degree one satisfying $d^2 = 0$, such that

$$d(gf) = g(df) + (-1)^p(dg)f$$

for all $g \in \mathrm{Hom}_{\mathbf{A}}(a, b)$ and all $f \in \mathrm{Hom}_{\mathbf{A}}(b, c)_p$ for $p \in \mathbb{Z}$. In this paper we will be interested in the category DG_n , which has as objects those DG categories with precisely n objects. If $\mathbf{A}, \mathbf{B} \in \mathrm{DG}_n$, recall that a DG functor $F: \mathbf{A} \rightarrow \mathbf{B}$ is a graded functor such that $F(df) = d(Ff)$ for all morphisms f . A *quasi-isomorphism* $F: \mathbf{A} \rightarrow \mathbf{B}$ is a DG functor inducing a bijection on objects, and quasi-isomorphisms $\mathrm{Hom}_{\mathbf{A}}(a_1, a_2) \rightarrow \mathrm{Hom}_{\mathbf{B}}(Fa_1, Fa_2)$ for all $a_1, a_2 \in \mathbf{A}$. Two categories $\mathbf{A}, \mathbf{B} \in \mathrm{DG}_n$ are called *quasi-isomorphic* if they are connected through a finite, non-directed chain of quasi-isomorphisms.

Notation 2.5. Suppose that $\mathbf{A} \in \mathrm{DG}_n$, and $(\Gamma, \mathbf{n}) \in \mathbf{Art}_n$, and recall from 2.3 that $\mathbf{n}_{ij} := e_i \mathbf{n} e_j$. Define $\mathbf{A} \underline{\otimes} \mathbf{n} := \bigoplus_{i,j=1}^n (\mathbf{A} \underline{\otimes} \mathbf{n})_{ij}$, where

$$(\mathbf{A} \underline{\otimes} \mathbf{n})_{ij} := \mathrm{Hom}_{\mathbf{A}}(i, j) \otimes_{\mathbb{C}} \mathbf{n}_{ij}.$$

Observe that $\mathbf{A} \underline{\otimes} \mathbf{n}$ has the natural structure of an object in DG_n (but with no units) with differential $d(a \otimes x) := d(a) \otimes x$. Thus we may consider $\mathbf{A} \underline{\otimes} \mathbf{n}$ as a DGLA, with bracket

$$[a \otimes x, b \otimes y] := ab \otimes xy - (-1)^{\deg(a) \deg(b)} ba \otimes yx$$

for homogeneous $a, b \in \mathbf{A}$.

Since $(\Gamma, \mathbf{n}) \in \mathbf{Art}_n$, by definition $\mathbf{n}^r = 0$ for some $r \geq 1$, hence $\mathbf{A} \underline{\otimes} \mathbf{n}$ is a nilpotent DGLA. This being the case, we can consider the standard Maurer–Cartan formulation to obtain a deformation functor.

Definition 2.6. Given $(\mathbf{A}, d) \in \mathrm{DG}_n$, the associated DG deformation functor

$$\mathrm{Def}^{\mathbf{A}}: \mathbf{Art}_n \rightarrow \mathbf{Sets}$$

is defined by sending

$$(\Gamma, \mathbf{n}) \mapsto \left\{ \xi \in \mathbf{A}^1 \underline{\otimes} \mathbf{n} \mid d(\xi) + \frac{1}{2}[\xi, \xi] = 0 \right\} / \sim$$

where as usual the equivalence relation \sim is induced by the gauge action. Explicitly, two elements $\xi_1, \xi_2 \in \mathbf{A}^1 \underline{\otimes} \mathbf{n}$ are said to be gauge equivalent if there exists $x \in \mathbf{A}^0 \underline{\otimes} \mathbf{n}$ such that

$$\xi_2 = e^x * \xi_1 := \xi_1 + \sum_{j=0}^{\infty} \frac{([x, -])^j}{(j+1)!} ([x, \xi_1] - d(x)).$$

The following is a mild extension of the well-known $n = 1$ case. The proof is very similar to the known $n = 1$ proofs (see e.g. [ELO, 8.1], [M99, 3.2], [GM, 2.4]), so we do not give it here.

Theorem 2.7. Suppose that $\mathbf{A} \rightarrow \mathbf{B}$ is a quasi-equivalence in DG_n . Then the deformation functors $\mathrm{Def}^{\mathbf{A}}$ and $\mathrm{Def}^{\mathbf{B}}$ are isomorphic.

2.3. Basic Results. Controlling noncommutative deformations of curves in the next section requires the following two preliminary results, and a corollary. All are well-known in the case $n = 1$.

To fix notation, suppose that \mathcal{A} is an abelian category and that c, d are two chain complexes with objects from \mathcal{A} . Set $\mathrm{Hom}_{\mathcal{A}}^{\mathrm{DG}}(x, y)$ to be the DG \mathbb{C} -module with

$$\mathrm{Hom}_{\mathcal{A}}^{\mathrm{DG}}(x, y)_t := \{(f_i)_{i \in \mathbb{Z}} \mid f_i: x_i \rightarrow y_{i+t}\}$$

and differential $\delta: f \mapsto fd_y - (-1)^{\deg(f)} d_x f$.

Now choose a family of objects $a_1, \dots, a_n \in \mathcal{A}$, an injective resolution $0 \rightarrow a_i \rightarrow I_\bullet^i$ for each a_i , and set $I := \bigoplus_{i=1}^n I_\bullet^i$. From this, we form $(\text{End}_{\mathcal{A}}^{\text{DG}}(I), \delta)$, considered naturally as an object of DG_n .

Lemma 2.8. *Suppose that \mathcal{A}, \mathcal{B} are abelian categories, and $F: \mathcal{A} \rightarrow \mathcal{B}$ is an additive functor with left adjoint L . Choose a family of objects $a_1, \dots, a_n \in \mathcal{A}$, and for each choose an injective resolution $0 \rightarrow a_i \rightarrow I_\bullet^i$. If*

- (1) L is exact,
- (2) $\mathbf{R}^t F(a_i) = 0$ for all $t > 0$ and all $i = 1, \dots, n$,
- (3) The counit $L \circ F \rightarrow \text{Id}$ is an isomorphism on each object a_i ,

then $\text{End}_{\mathcal{A}}^{\text{DG}}(\bigoplus I_\bullet^i)$ and $\text{End}_{\mathcal{B}}^{\text{DG}}(\bigoplus FI_\bullet^i)$ are quasi-isomorphic in DG_n .

Proof. Consider the obvious map

$$F: \text{End}_{\mathcal{A}}^{\text{DG}}\left(\bigoplus I_\bullet^i\right) \rightarrow \text{End}_{\mathcal{B}}^{\text{DG}}\left(\bigoplus FI_\bullet^i\right). \quad (2.A)$$

Condition (1) implies that F preserves injective objects, and (2) implies that F preserves the injective resolutions of the a_i , hence $0 \rightarrow Fa_i \rightarrow FI_\bullet^i$ are injective resolutions. Hence the cohomologies of the left hand side of (2.A) compute $\text{Ext}_{\mathcal{A}}^n(a_i, a_j)$, and the right hand side computes $\text{Ext}_{\mathcal{B}}^n(Fa_i, Fa_j)$. These are obviously the same, via F , since by (3)

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Hom}_{\mathcal{A}}(a_i, I_0^j) & \longrightarrow & \text{Hom}_{\mathcal{A}}(a_i, I_1^j) & \longrightarrow & \text{Hom}_{\mathcal{A}}(a_i, I_2^j) \longrightarrow \dots \\ & & \wr \downarrow & & \wr \downarrow & & \wr \downarrow \\ 0 & \longrightarrow & \text{Hom}_{\mathcal{A}}(LFa_i, I_0^j) & \longrightarrow & \text{Hom}_{\mathcal{A}}(LFa_i, I_1^j) & \longrightarrow & \text{Hom}_{\mathcal{A}}(LFa_i, I_2^j) \longrightarrow \dots \\ & & \wr \downarrow & & \wr \downarrow & & \wr \downarrow \\ 0 & \longrightarrow & \text{Hom}_{\mathcal{B}}(Fa_i, FI_0^j) & \longrightarrow & \text{Hom}_{\mathcal{B}}(Fa_i, FI_1^j) & \longrightarrow & \text{Hom}_{\mathcal{B}}(Fa_i, FI_2^j) \longrightarrow \dots \end{array}$$

commutes and all the vertical morphisms are isomorphisms. \square

Keeping the notation as above, for each of the objects $a_1, \dots, a_n \in \mathcal{A}$, choose a left resolution $Q_i^\bullet \rightarrow a_i \rightarrow 0$, where for now the Q 's are arbitrary. Set $Q := \bigoplus_{i=1}^n Q_i^\bullet$, and consider

$$\Delta := \begin{pmatrix} \text{End}_{\mathcal{A}}^{\text{DG}}(Q[1]) & \text{Hom}_{\mathcal{A}}^{\text{DG}}(Q[1], I) \\ 0 & \text{End}_{\mathcal{A}}^{\text{DG}}(I) \end{pmatrix}.$$

This can be viewed as an object in DG_n in the obvious way: the homomorphism space between object i and object j is

$$\begin{pmatrix} \text{End}_{\mathcal{A}}^{\text{DG}}(Q_i^\bullet[1]) & \text{Hom}_{\mathcal{A}}^{\text{DG}}(Q_i^\bullet[1], I_\bullet^j) \\ 0 & \text{End}_{\mathcal{A}}^{\text{DG}}(I_\bullet^j) \end{pmatrix},$$

with differential as in [ELO, §8]. There are natural projections $p_1: \Delta \rightarrow \text{End}_{\mathcal{A}}^{\text{DG}}(Q[1])$ and $p_2: \Delta \rightarrow \text{End}_{\mathcal{A}}^{\text{DG}}(I)$ in DG_n , and splicing the left resolutions with the right resolutions gives an exact complex $Q[1] \rightarrow I$.

Proposition 2.9 (Keller). *Suppose that \mathcal{A} is an abelian category, and choose a family of objects $a_1, \dots, a_n \in \mathcal{A}$. With notation as above,*

- (1) *The projection p_2 is a quasi-isomorphism in DG_n .*
- (2) *If $\text{Hom}_{\mathcal{A}}^{\text{DG}}(Q[1], Q[1] \rightarrow I)$ is exact, then p_1 is a quasi-isomorphism in DG_n .*

In particular, provided that $\text{Hom}_{\mathcal{A}}^{\text{DG}}(Q[1], Q[1] \rightarrow I)$ is exact, $\text{End}_{\mathcal{A}}^{\text{DG}}(Q)$ and $\text{End}_{\mathcal{A}}^{\text{DG}}(I)$ are quasi-isomorphic in DG_n .

Proof. As above, the complex $Q[1] \rightarrow I$ is exact.

- (1) By construction of the upper triangular Δ , as in [ELO, §8]

$$\text{Ker } p_2 = \text{Hom}_{\mathcal{A}}^{\text{DG}}(I, Q[1] \rightarrow I).$$

Since I is h -injective, it follows that $\text{Ker } p_2$ is exact, and thus p_2 is a quasi-isomorphism.

- (2) Again by construction, $\text{Ker } p_1 = \text{Hom}_{\mathcal{A}}^{\text{DG}}(Q[1], Q[1] \rightarrow I)$, and so if by assumption this is exact, p_1 is a quasi-isomorphism.

The final statement follows from (1) and (2), since clearly $\text{End}_{\mathcal{A}}^{\text{DG}}(Q) \cong \text{End}_{\mathcal{A}}^{\text{DG}}(Q[1])$. \square

The following is now a direct consequence of [S08, 2.8], and says that the naive deformations and the DG deformations are the same when we deform distinct simples.

Corollary 2.10. *Suppose that Λ is a \mathbb{C} -algebra, and that $\mathcal{S} = \{S_1, \dots, S_n\} \subseteq \text{Mod } \Lambda$ are simple and distinct. Choose injective resolutions $0 \rightarrow S_i \rightarrow I_\bullet^i$ and set $A := \text{End}_\Lambda^{\text{DG}}(I) \in \text{DG}_n$ as above. Then $\text{Def}^A \cong \text{Def}_\Lambda^{\mathcal{S}}$.*

Proof. Under the assumption that the S_i are distinct simples, Segal [S08, 2.6, 2.8] shows that $\text{Def}_\Lambda^{\mathcal{S}}$ is isomorphic to the DG deformation functor associated to the bar resolutions of the simples. Since projective resolutions are h -projective, the conditions of 2.9(2) are satisfied, so the bar resolution DGA is quasi-isomorphic in DG_n to A . The result then follows from 2.7. \square

3. DEFORMATIONS OF REDUCED FIBRES

In this section, in the setting of contractions with at most one-dimensional fibres, we show that the functor of simultaneous noncommutative deformations of the reduced fibre is prorepresented by a naturally defined algebra Λ_{con} . This algebra is a factor of one obtained by tilting, and this extra control over the prorepresenting object allows us in §4 to prove that noncommutative deformations recover the contracted locus.

3.1. Setup. This subsection fixes notation. Throughout the paper, we will refer to the three setups in 3.1, 3.2 and 3.4 below.

Setup 3.1. (Global) Suppose that $f: X \rightarrow X_{\text{con}}$ is a projective birational morphism between noetherian integral normal \mathbb{C} -schemes, with $\mathbf{R}f_*\mathcal{O}_X = \mathcal{O}_{X_{\text{con}}}$, such that the fibres are at most one-dimensional. Throughout, we write L for the locus of (not necessarily closed) points of X_{con} above which f is not an isomorphism.

We make no assumptions on the singularities of X . Next, for any closed point $p \in L$, we pick an affine neighbourhood $\text{Spec } R$ in X_{con} containing p , and after base change consider the following Zariski local setup.

Setup 3.2. (Zariski local) Suppose that $f: U \rightarrow \text{Spec } R$ is a projective birational morphism between noetherian integral normal \mathbb{C} -schemes, with $\mathbf{R}f_*\mathcal{O}_U = \mathcal{O}_R$, such that the fibres are at most one-dimensional.

In dimension 3, an easy example is the blowup of \mathbb{A}^3 at the ideal (x, y) , but the setup also includes arbitrary flips and flops of multiple curves, as well as divisorial contractions to curves. We make no assumptions on the singularities of U .

With the assumptions in 3.2, it is well-known [V04, 3.2.8] that there is a bundle $\mathcal{V} := \mathcal{O}_U \oplus \mathcal{N}$ inducing a derived equivalence

$$\text{D}^b(\text{coh } U) \xrightarrow[\sim]{\mathbf{R}\text{Hom}_U(\mathcal{V}, -)} \text{D}^b(\text{mod } \text{End}_U(\mathcal{V})). \quad (3.A)$$

Throughout we set

$$\Lambda := \text{End}_U(\mathcal{V}) = \text{End}_U(\mathcal{O}_U \oplus \mathcal{N}),$$

and recall from the conventions in §2.2 that if $\mathcal{F}, \mathcal{G} \in \text{coh } U$ where \mathcal{F} is a summand of \mathcal{G} , then we define the ideal $[\mathcal{F}]$ to be the two-sided ideal of $\text{End}_U(\mathcal{G})$ consisting of all morphisms factoring through $\text{add } \mathcal{F}$.

The following is similar to [DW1, 2.12], but the definition is now more subtle since in general $\text{End}_U(\mathcal{V}) \not\cong \text{End}_R(f_*\mathcal{V})$, whereas there is such an isomorphism in the setting of [DW1]. The upshot is that we must work on U , and not $\text{Spec } R$.

Definition 3.3. *With notation as above, we define the contraction algebra associated to Λ to be $\Lambda_{\text{con}} := \text{End}_U(\mathcal{O}_U \oplus \mathcal{N})/[\mathcal{O}_U]$.*

The algebra Λ_{con} defined above depends on Λ and thus the choice of derived equivalence (3.A), but this is accounted for in the formal fibre setting below, after passing through morita equivalences. Also, we remark that since $\mathcal{N} \notin \text{add } \mathcal{O}_U$ (else f is an isomorphism, e.g. by 4.5), the contraction algebra Λ_{con} is necessarily non-zero.

To obtain well-defined invariants that do not depend on choices, and also to relate to the deformation theory in the following §3.2, we now pass to the formal fibre.

Setup 3.4. (Complete local) Suppose that $f: \mathfrak{U} \rightarrow \operatorname{Spec} \mathfrak{R}$ is a projective birational morphism between noetherian integral normal \mathbb{C} -schemes, with $\mathbf{R}f_* \mathcal{O}_{\mathfrak{U}} = \mathcal{O}_{\mathfrak{R}}$, where \mathfrak{R} is complete local and the fibres of f are at most one-dimensional.

After passing to this formal fibre, the Zariski local derived equivalence has a particularly nice form, which we now briefly review. Fix a closed point $\mathfrak{m} \in L$, and write $\mathfrak{R} := \widehat{R}$. The above derived equivalence (3.A) induces an equivalence

$$\mathrm{D}^b(\operatorname{coh} \mathfrak{U}) \xrightarrow[\sim]{\mathbf{R}\operatorname{Hom}_{\mathfrak{U}}(\widehat{\mathcal{V}}, -)} \mathrm{D}^b(\operatorname{mod} \widehat{\Lambda}),$$

and this can be described much more explicitly. We let $C = \pi^{-1}(\mathfrak{m})$ where \mathfrak{m} is the unique closed point of $\operatorname{Spec} \mathfrak{R}$, then giving C the reduced scheme structure, we can write $C^{\operatorname{red}} = \bigcup_{i=1}^n C_i$ with each $C_i \cong \mathbb{P}^1$. Let \mathcal{L}_i denote the line bundle on \mathfrak{U} such that $\mathcal{L}_i \cdot C_j = \delta_{ij}$. If the multiplicity of C_i is equal to one, set $\mathcal{M}_i := \mathcal{L}_i$, else define \mathcal{M}_i to be given by the maximal extension

$$0 \rightarrow \mathcal{O}_{\mathfrak{U}}^{\oplus(r-1)} \rightarrow \mathcal{M}_i \rightarrow \mathcal{L}_i \rightarrow 0$$

associated to a minimal set of $r-1$ generators of $H^1(\mathfrak{U}, \mathcal{L}_i^*)$ as an \mathfrak{R} -module [V04, 3.5.4]. Then $\mathcal{O}_{\mathfrak{U}} \oplus \bigoplus_{i=1}^n \mathcal{M}_i^*$ is a tilting bundle on \mathfrak{U} [V04, 3.5.5] generating ${}^0\mathcal{P}er(\mathfrak{U})$. By [V04, 3.5.5] we can write

$$\mathcal{O}_{\mathfrak{U}} \oplus \widehat{\mathcal{N}} \cong \mathcal{O}_{\mathfrak{U}}^{\oplus a_0} \oplus \bigoplus_{i=1}^n (\mathcal{M}_i^*)^{\oplus a_i}$$

for some $a_i \in \mathbb{N}$ and so consequently $\widehat{\Lambda} \cong \operatorname{End}_{\mathfrak{U}}(\mathcal{O}_{\mathfrak{U}}^{\oplus a_0} \oplus \bigoplus_{i=1}^n (\mathcal{M}_i^*)^{\oplus a_i})$.

Definition 3.5. We write $\mathcal{M}^* := \bigoplus_{i=1}^n \mathcal{M}_i^*$ and define

$$\mathbf{A} := \operatorname{End}_{\mathfrak{U}}(\mathcal{O}_{\mathfrak{U}} \oplus \mathcal{M}^*),$$

which is the basic algebra morita equivalent to $\widehat{\Lambda}$. From this, we define the contraction algebra associated to f to be

$$\mathbf{A}_{\operatorname{con}} := \operatorname{End}_{\mathfrak{U}}(\mathcal{O}_{\mathfrak{U}} \oplus \mathcal{M}^*) / [\mathcal{O}_{\mathfrak{U}}] \cong \operatorname{End}_{\mathfrak{U}}(\mathcal{M}^*) / [\mathcal{O}_{\mathfrak{U}}],$$

and we define the fibre algebra associated to f to be

$$\mathbf{A}_{\operatorname{fib}} := \operatorname{End}_{\mathfrak{U}}(\mathcal{O}_{\mathfrak{U}} \oplus \mathcal{M}^*) / [\mathcal{M}^*] \cong \operatorname{End}_{\mathfrak{U}}(\mathcal{O}_{\mathfrak{U}}) / [\mathcal{M}^*].$$

Remark 3.6. Since $\operatorname{End}_{\mathfrak{U}}(\mathcal{O}_{\mathfrak{U}}) \cong \mathfrak{R}$, it follows from the definition that $\mathbf{A}_{\operatorname{fib}}$ is always commutative, although $\mathbf{A}_{\operatorname{con}}$ need not be.

To establish various homological properties we will need to pass through morita equivalences between the algebras \mathbf{A} and $\widehat{\Lambda}$, and between the algebras $\mathbf{A}_{\operatorname{con}}$ and $\widehat{\Lambda}_{\operatorname{con}}$. Here we describe these equivalences, mainly to fix notation. In analogy with [DW1, §5.3] throughout we write

$$\mathcal{Y} := \mathcal{O}_{\mathfrak{U}} \oplus \bigoplus \mathcal{M}_i^*, \quad \mathcal{Z} := \mathcal{O}_{\mathfrak{U}}^{\oplus a_0} \oplus \bigoplus (\mathcal{M}_i^*)^{\oplus a_i},$$

so that $\mathbf{A} = \operatorname{End}_{\mathfrak{U}}(\mathcal{Y})$ and $\widehat{\Lambda} = \operatorname{End}_{\mathfrak{U}}(\mathcal{Z})$. Writing

$$P := \operatorname{Hom}_{\mathfrak{U}}(\mathcal{Y}, \mathcal{Z}), \quad Q := \operatorname{Hom}_{\mathfrak{U}}(\mathcal{Z}, \mathcal{Y})$$

it is clear that both P and Q have the structure of bimodules, namely ${}_{\widehat{\Lambda}}P_{\mathbf{A}}$ and ${}_{\mathbf{A}}Q_{\widehat{\Lambda}}$. It is easy to see that P is a progenerator, and that this induces the following result.

Lemma 3.7. *With notation as above, there is a morita equivalence*

$$\operatorname{mod} \mathbf{A} \xrightleftharpoons[\operatorname{Hom}_{\widehat{\Lambda}}(Q, -) = - \otimes_{\widehat{\Lambda}} P]{\operatorname{Hom}_{\mathbf{A}}(P, -) = - \otimes_{\mathbf{A}} Q} \operatorname{mod} \widehat{\Lambda}, \quad (3.B)$$

and a morita equivalence

$$\operatorname{mod} \mathbf{A}_{\operatorname{con}} \xrightleftharpoons[\operatorname{Hom}_{\widehat{\Lambda}_{\operatorname{con}}}(\mathbb{F}\mathbf{A}_{\operatorname{con}}, -)]{- \otimes_{\mathbf{A}_{\operatorname{con}}} \mathbb{F}\mathbf{A}_{\operatorname{con}}} \operatorname{mod} \widehat{\Lambda}_{\operatorname{con}}. \quad (3.C)$$

3.2. Deformations and Contraction Algebras. In this subsection we will prove that the contraction algebra A_{con} prorepresents various natural deformation functors. We will translate deformation problems across a variety of different functors, so we now set notation, as in [DW3, (2.G)].

Notation 3.8. We pick a closed point $\mathfrak{m} \in L$, and as above consider $C := f^{-1}(\mathfrak{m})$. We write $C^{\text{red}} = \bigcup_{i=1}^n C_i$ with each $C_i \cong \mathbb{P}^1$, and put T_i for the simple Λ -modules corresponding to the (perverse) sheaves $\mathcal{O}_{C_i}(-1)$ across the derived equivalence (3.A). Further, we denote the simple A -modules by $S_i := \mathbb{F}^{-1}\widehat{T}_i$, so that

$$\begin{array}{ccccc} \text{D}^b(\text{Qcoh } U) & \xrightleftharpoons[\text{-} \otimes_{\Lambda}^{\mathbf{L}} \mathcal{V}]{\mathbf{R}\text{Hom}_U(\mathcal{V}, -)} & \text{D}^b(\text{Mod } \Lambda) & \xrightleftharpoons[\text{res} \circ \mathbb{F}]{\mathbb{F}^{-1} \circ (-)} & \text{D}^b(\text{Mod } A) \\ \mathcal{O}_{C_i}(-1) & \longleftrightarrow & T_i & \longrightarrow & S_i \end{array}$$

For the remainder of this subsection, we will use the following simplified notation:

- (1) $\mathcal{D}ef_X$ for $\mathcal{D}ef^{A_1}$, where $A_1 \in \text{DG}_n$ is obtained from the injective resolutions of the $\mathcal{O}_{C_i}(-1) \in \text{coh } X$.
- (2) $\mathcal{D}ef_U$ for $\mathcal{D}ef^{A_2}$, where $A_2 \in \text{DG}_n$ is obtained from the injective resolutions of the $\mathcal{O}_{C_i}(-1) \in \text{coh } U$.
- (3) $\mathcal{D}ef_{\Lambda}$ for $\mathcal{D}ef^{A_3}$, where $A_3 \in \text{DG}_n$ is obtained from the injective resolutions of the $T_i \in \text{mod } \Lambda$.
- (4) $\mathcal{D}ef_A$ for $\mathcal{D}ef^{A_4}$, where $A_4 \in \text{DG}_n$ is obtained from the injective resolutions of the $S_i \in \text{mod } A$.

The next result is now an easy corollary of the DG results in §2.3, vastly simplifying [DW1, 2.6, 2.8, 2.19].

Theorem 3.9. *With the global setup of 3.1, and notation in 3.8,*

- (1) $\mathcal{D}ef_X \cong \mathcal{D}ef_U$.
- (2) $\mathcal{D}ef_U \cong \mathcal{D}ef_{\Lambda}$.
- (3) $\mathcal{D}ef_{\Lambda} \cong \mathcal{D}ef_A$.
- (4) $\mathcal{D}ef_A \cong \text{Hom}_{\text{pArt}_n}(A_{\text{con}}, -)$.

In particular, all the deformation functors above are prorepresented by A_{con} .

Proof. By 2.7, we just need to show that the DGAs are quasi-isomorphic.

(1) Consider $F := i_*: \text{Qcoh } U \rightarrow \text{Qcoh } X$, with exact left adjoint $L = i^*$. Since C_i is closed, $\mathbf{R}i_*\mathcal{O}_{C_i}(-1) = i_*\mathcal{O}_{C_i}(-1)$ for each $i = 1, \dots, n$. Further, the counit $L \circ F \rightarrow \text{Id}$ is an isomorphism for all sheaves, in particular for the sheaves $\mathcal{O}_{C_i}(-1)$. Hence the result follows from 2.8.

(2) Set $F := \text{Hom}_U(\mathcal{V}, -): \text{Qcoh } U \rightarrow \text{Mod } \Lambda$, with (non-exact) left adjoint $L := \mathcal{V} \otimes_{\Lambda} -$. To establish the claim we will first use 2.9, so set $E_i := \mathcal{O}_{C_i}(-1)$, and for each $i = 1, \dots, n$ choose an injective resolution $0 \rightarrow E_i \rightarrow I_{\bullet}^i$. On the other hand, choose a projective resolution $P_i^{\bullet} \rightarrow T_i \rightarrow 0$ of each of the T_i . Since the sheaf E_i corresponds to the module T_i across the equivalence (3.A), it follows that $\mathcal{V} \otimes_{\Lambda}^{\mathbf{L}} T_i \cong E_i$. In particular all higher cohomology groups vanish, and so $LP_i^{\bullet} \rightarrow E_i \rightarrow 0$ is exact, giving a locally free resolution of E_i .

To match the notation of 2.9, set $Q_i^{\bullet} := LP_i^{\bullet}$, and write $P = \bigoplus_{i=1}^n P_i^{\bullet}$, $Q = \bigoplus_{i=1}^n Q_i^{\bullet}$, and $I = \bigoplus_{i=1}^n I_{\bullet}^i$. By adjunction

$$\text{Hom}_U^{\text{DG}}(LP[1], LP[1] \rightarrow I) \cong \text{Hom}_{\Lambda}^{\text{DG}}(P[1], P[1] \rightarrow FI). \quad (3.D)$$

Since P is h -projective, and $P[1] \rightarrow FI$ is acyclic, (3.D) is exact. By 2.9, this then establishes that $\text{End}_U^{\text{DG}}(I)$ and $\text{End}_U^{\text{DG}}(Q)$ are quasi-isomorphic.

Now it is clear that

$$\text{End}_{\Lambda}^{\text{DG}}(P) \rightarrow \text{End}_U^{\text{DG}}(LP) = \text{End}_U^{\text{DG}}(Q)$$

is a quasi-isomorphism, and so combining we see that $\text{End}_{\Lambda}^{\text{DG}}(P)$ is quasi-isomorphic to $\text{End}_U^{\text{DG}}(I)$. Finally, choose an injective resolution $0 \rightarrow T_i \rightarrow J_{\bullet}^i$ of each T_i , and set $J := \bigoplus_{i=1}^n J_{\bullet}^i$. It is well known (again using 2.9), that $\text{End}_{\Lambda}^{\text{DG}}(J)$ is quasi-isomorphic to $\text{End}_{\Lambda}^{\text{DG}}(P)$, and thus by the above to $\text{End}_U^{\text{DG}}(I)$.

(3) Consider $F := \mathbb{F}^{-1} \circ \widehat{(-)}: \text{Mod } \Lambda \rightarrow \text{Mod } A$, which has exact left adjoint $L = \text{res} \circ \mathbb{F}$. Being the composition of exact functors, F is also exact, so $\mathbf{R}^t F(S_i) = 0$ for all $t > 0$ and for all $i = 1, \dots, n$. Since \mathbb{F} is an equivalence, and since the adjunction $\text{res} \dashv \widehat{(-)}$ restricts to an equivalence between finite length Λ -modules supported at \mathfrak{m} and finite length $\widehat{\Lambda}$ -modules, it follows that the counit $L \circ F \rightarrow \text{Id}$ is an isomorphism on the objects S_i . Hence the statement follows from 2.8.

(4) By 2.10, the DG deformation functor Def_A is isomorphic to the naive deformation functor in 2.4. Thus, since each S_i is one-dimensional, using the standard argument (see e.g. [DW1, 3.1]) it follows that

$$\begin{aligned} \text{Def}_A(\Gamma) &= \left\{ \begin{array}{l} \bullet \text{ A left } A\text{-module structure on } M = (S_i \otimes_{\mathbb{C}} \Gamma_{ij}) \text{ such that} \\ \quad (S_i \otimes_{\mathbb{C}} \Gamma_{ij}) \text{ becomes a } A\text{-}\Gamma \text{ bimodule.} \\ \bullet \text{ A collection of isomorphisms } \delta_i: M \otimes_{\Gamma} (\Gamma/\mathfrak{m})e_i \xrightarrow{\sim} S_i \end{array} \right\} / \sim \\ &= \left\{ \begin{array}{l} \bullet \text{ A left } A\text{-module structure on } \Gamma \text{ such that } \Gamma \in \text{Mod } A \otimes \Gamma^{\text{op}}. \\ \bullet \text{ A collection of isomorphisms } \delta_i: \Gamma \otimes_{\Gamma} (\Gamma/\mathfrak{m})e_i \xrightarrow{\sim} S_i \end{array} \right\} / \sim \\ &= \text{Hom}_{\text{Alg}_n}(A_{\text{con}}, \Gamma). \end{aligned}$$

It remains to show that $A_{\text{con}} \in \mathbf{pArt}_n$, but this holds since \mathfrak{A} , thus A , and thus A_{con} , are complete with respect to their augmentation ideals. \square

Remark 3.10. The above proof of 3.9(2) establishes that we can compute Def_U using the DGA associated to the specific locally free resolutions $\mathcal{V} \otimes_{\Lambda} P_i^{\bullet}$ of the E_i . It is rare to be able to compute the deformations of a sheaf using the DGA of a locally free resolution, and essentially it is this extra control of the deformation theory that allows us to prove the contraction theorem in 4.8 below. Note that even taking the DGA of a different locally free resolution of the E_i may give a different deformation functor.

4. THE CONTRACTION THEOREM

In the global setup of 3.1, by 3.9 it follows that simultaneous noncommutative deformations of the reduced fibre is prorepsented by A_{con} , namely

$$\text{Def}_X \cong \text{Hom}_{\mathbf{pArt}_n}(A_{\text{con}}, -).$$

Further, by 3.7, A_{con} is morita equivalent to $\widehat{\Lambda}_{\text{con}}$. We are thus motivated to control A_{con} (and Λ_{con}), and we now do this in a sequence of reduction steps, culminating in the contraction theorem of §4.2.

4.1. Behaviour of Λ_{con} under Global Sections and Base Change. We revert to the Zariski local setup of 3.2. The following lemma is important: it is very well-known if f is an isomorphism in codimension two [V04, 4.2.1], however in our more general setting care is required.

Lemma 4.1. *With the setup as in 3.2, $\text{End}_U(\mathcal{V}^*) \cong \text{End}_R(f_*(\mathcal{V}^*))$ so*

$$\Lambda \cong \text{End}_R(R \oplus f_*(\mathcal{N}^*))^{\text{op}} \quad \text{and} \quad \Lambda_{\text{con}} \cong (\text{End}_R(R \oplus f_*(\mathcal{N}^*)) / [f_*(\mathcal{N}^*)])^{\text{op}}.$$

This allows us to reduce many problems to the base $\text{Spec } R$. Indeed $\text{End}_U(\mathcal{V}) \not\cong \text{End}_R(f_*\mathcal{V})$ in general, so the statement and proof of 4.1 is a little subtle. Since \mathcal{V}^* is generated by global sections and is tilting, the proof of 4.1 follows immediately from 4.3 below. This requires the following easy well-known lemma.

Lemma 4.2. *Assume that $f: Y \rightarrow \text{Spec } S$ is a proper birational map between integral schemes, where S is normal.*

- (1) *Suppose that $\mathcal{F} \in \text{coh } Y$ is generated by global sections. Then we can find a morphism $\mathcal{O}_Y^{\oplus a} \rightarrow \mathcal{F}$ such that $S^{\oplus a} \rightarrow f_*\mathcal{F}$.*
- (2) *Suppose that $\mathcal{W}_1, \mathcal{W}_2$ are finite rank vector bundles on Y . Then*

$$\text{Hom}_Y(\mathcal{W}_1, \mathcal{W}_2) \hookrightarrow \text{Hom}_S(f_*\mathcal{W}_1, f_*\mathcal{W}_2).$$

Proof. (1) This is a basic consequence of Zariski's main theorem, together with the fact that f_* preserves coherence.

(2) Since $\text{Hom}_Y(\mathcal{W}_1, \mathcal{W}_2) = H^0(\mathcal{W}_1^* \otimes \mathcal{W}_2)$ and $\mathcal{W}_1^* \otimes \mathcal{W}_2$ is a vector bundle, by integrality $\text{Hom}_Y(\mathcal{W}_1, \mathcal{W}_2)$ is a torsion-free S -module. The statement follows. \square

Proposition 4.3. *Assume that $f: Y \rightarrow \operatorname{Spec} S$ is a proper birational map between integral schemes, where S is normal. If \mathcal{W} is a vector bundle on Y of finite rank, which is generated by global sections, such that $\operatorname{Ext}_Y^1(\mathcal{W}, \mathcal{W}) = 0$, then $\operatorname{End}_Y(\mathcal{W}) \cong \operatorname{End}_S(f_*\mathcal{W})$.*

Proof. By 4.2(1) there is a short exact sequence

$$0 \rightarrow \mathcal{K} \rightarrow \mathcal{O}_Y^{\oplus a} \rightarrow \mathcal{W} \rightarrow 0 \quad (4.A)$$

such that

$$0 \rightarrow f_*\mathcal{K} \rightarrow S^{\oplus a} \rightarrow f_*\mathcal{W} \rightarrow 0 \quad (4.B)$$

is exact. Applying $\operatorname{Hom}_Y(-, \mathcal{W})$ to (4.A) and applying $\operatorname{Hom}_S(-, f_*\mathcal{W})$ to (4.B), we have an exact commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \operatorname{Hom}_Y(\mathcal{W}, \mathcal{W}) & \longrightarrow & \operatorname{Hom}_Y(\mathcal{O}_Y^{\oplus a}, \mathcal{W}) & \longrightarrow & \operatorname{Hom}_Y(\mathcal{K}, \mathcal{W}) \rightarrow \operatorname{Ext}_Y^1(\mathcal{W}, \mathcal{W}) = 0 \\ & & \downarrow \alpha & & \parallel & & \downarrow \beta \\ 0 & \longrightarrow & \operatorname{Hom}_S(f_*\mathcal{W}, f_*\mathcal{W}) & \longrightarrow & \operatorname{Hom}_S(S^{\oplus a}, f_*\mathcal{W}) & \longrightarrow & \operatorname{Hom}_S(f_*\mathcal{K}, f_*\mathcal{W}) \end{array}$$

where both α and β are injective by 4.2(2). By the snake lemma, α is also surjective. \square

With the setup in 3.2, both Λ and $\Lambda_{\operatorname{con}}$ have the structure of an R -module. For our purposes later, we need to control this under flat base change. In what follows, for $\mathfrak{p} \in \operatorname{Spec} R$ we consider the base change squares

$$\begin{array}{ccccc} \mathfrak{U}_{\mathfrak{p}} & \xrightarrow{m} & U_{\mathfrak{p}} & \xrightarrow{k} & U \\ \theta \downarrow & & \varphi \downarrow & & \downarrow f \\ \operatorname{Spec} \mathfrak{R}_{\mathfrak{p}} & \xrightarrow{l} & \operatorname{Spec} R_{\mathfrak{p}} & \xrightarrow{j} & \operatorname{Spec} R \end{array}$$

where $\mathfrak{R}_{\mathfrak{p}}$ denotes the completion of $R_{\mathfrak{p}}$ at its unique maximal ideal.

Proposition 4.4. *With the setup as in 3.2,*

- (1) $U_{\mathfrak{p}}$ is derived equivalent to $\Lambda_{\mathfrak{p}}$ via the tilting bundle $k^*\mathcal{V} = \mathcal{O}_{U_{\mathfrak{p}}} \oplus k^*\mathcal{N}$.
- (2) $\mathfrak{U}_{\mathfrak{p}}$ is derived equivalent to $\widehat{\Lambda}_{\mathfrak{p}}$ via the tilting bundle $m^*k^*\mathcal{V} = \mathcal{O}_{\mathfrak{U}_{\mathfrak{p}}} \oplus m^*k^*\mathcal{N}$.
- (3) $(\Lambda_{\operatorname{con}})_{\mathfrak{p}} \cong (\Lambda_{\mathfrak{p}})_{\operatorname{con}}$.
- (4) $(\Lambda_{\operatorname{con}})_{\mathfrak{p}} \otimes_{R_{\mathfrak{p}}} \mathfrak{R}_{\mathfrak{p}} \cong (\widehat{\Lambda}_{\mathfrak{p}})_{\operatorname{con}}$.

Proof. All statements are elementary, but we give the proof for completeness.

(1) U is derived equivalent to Λ via the tilting bundle $\mathcal{V} := \mathcal{O}_U \oplus \mathcal{N}$. It is well-known that this implies $U_{\mathfrak{p}}$ is derived equivalent to $\Lambda_{\mathfrak{p}}$ via the tilting bundle $k^*\mathcal{V} = \mathcal{O}_{U_{\mathfrak{p}}} \oplus k^*\mathcal{N}$. For example, a proof of the Ext vanishing together with the fact that $\operatorname{End}_{U_{\mathfrak{p}}}(k^*\mathcal{V}) \cong \Lambda_{\mathfrak{p}}$ can be found in [IW1, 4.3(2)]. For generation, first observe that j is an affine morphism, hence so is k . Then $\mathbf{R}\operatorname{Hom}_{U_{\mathfrak{p}}}(k^*\mathcal{V}, x) = 0$ implies, by adjunction, that $\mathbf{R}\operatorname{Hom}_U(\mathcal{V}, k_*x) = 0$. Since \mathcal{V} generates, $k_*x = 0$ and so since k is affine $x = 0$.

(2) The proof is identical to (1).

(3) Since f is proper, the category $\operatorname{coh} U$ is R -linear. In particular $\operatorname{Hom}_U(\mathcal{O}_U, \mathcal{V})$ is a finitely generated $\operatorname{End}_U(\mathcal{O}_U)$ -module, so we can find a surjection

$$\operatorname{Hom}_U(\mathcal{O}_U, \mathcal{O}_U)^{\oplus a} \rightarrow \operatorname{Hom}_U(\mathcal{O}_U, \mathcal{V}) \rightarrow 0. \quad (4.C)$$

Tracking the images of the identities on the left hand side under the above map gives elements $g_1, \dots, g_a \in \operatorname{Hom}_U(\mathcal{O}_U, \mathcal{V})$, and so we may use these to form a natural morphism

$$\mathcal{O}_U^{\oplus a} \xrightarrow{h} \mathcal{V} \quad (4.D)$$

such that applying $\operatorname{Hom}_U(\mathcal{O}_U, -)$ to (4.D) gives (4.C). By definition, this means that h is an add \mathcal{O}_U -approximation of \mathcal{V} . Hence applying $\operatorname{Hom}_U(\mathcal{V}, -)$ to (4.D) yields an exact sequence

$$\operatorname{Hom}_U(\mathcal{V}, \mathcal{O}_U)^{\oplus a} \rightarrow \operatorname{Hom}_U(\mathcal{V}, \mathcal{V}) \rightarrow \Lambda_{\operatorname{con}} \rightarrow 0. \quad (4.E)$$

Interpreting $\mathrm{Hom}_U(-, -) = f_* \mathcal{H}om(-, -)$, applying the exact functor $j^*(-) = (-)_{\mathfrak{p}}$ to (4.E) and using flat base change gives the exact sequence

$$\varphi_* k^* \mathcal{H}om_U(\mathcal{V}, \mathcal{O}_U)^{\oplus a} \rightarrow \varphi_* k^* \mathcal{H}om_U(\mathcal{V}, \mathcal{V}) \rightarrow (\Lambda_{\mathrm{con}})_{\mathfrak{p}} \rightarrow 0.$$

Since \mathcal{V} is coherent we may move the k^* inside $\mathcal{H}om$, and so the above is simply

$$\mathrm{Hom}_{U_{\mathfrak{p}}}(k^* \mathcal{V}, \mathcal{O}_{U_{\mathfrak{p}}})^{\oplus a} \rightarrow \mathrm{Hom}_{U_{\mathfrak{p}}}(k^* \mathcal{V}, k^* \mathcal{V}) \rightarrow (\Lambda_{\mathrm{con}})_{\mathfrak{p}} \rightarrow 0. \quad (4.F)$$

But on the other hand applying the exact functor k^* to (4.D) gives a morphism

$$\mathcal{O}_{U_{\mathfrak{p}}}^{\oplus a} \xrightarrow{k^*(h)} k^* \mathcal{V} \quad (4.G)$$

and further applying j^* to (4.C) and using flat base change shows that

$$\mathrm{Hom}_{U_{\mathfrak{p}}}(\mathcal{O}_{U_{\mathfrak{p}}}, \mathcal{O}_{U_{\mathfrak{p}}})^{\oplus a} \rightarrow \mathrm{Hom}_{U_{\mathfrak{p}}}(\mathcal{O}_{U_{\mathfrak{p}}}, k^* \mathcal{V}) \rightarrow 0$$

is exact. Hence $k^*(h)$ is an add $\mathcal{O}_{U_{\mathfrak{p}}}$ -approximation, and so applying $\mathrm{Hom}_{U_{\mathfrak{p}}}(k^* \mathcal{V}, -)$ to (4.G) gives an exact sequence

$$\mathrm{Hom}_{U_{\mathfrak{p}}}(k^* \mathcal{V}, \mathcal{O}_{U_{\mathfrak{p}}})^{\oplus a} \rightarrow \mathrm{Hom}_{U_{\mathfrak{p}}}(k^* \mathcal{V}, k^* \mathcal{V}) \rightarrow (\Lambda_{\mathfrak{p}})_{\mathrm{con}} \rightarrow 0. \quad (4.H)$$

Combining (4.F) and (4.H) shows that $(\Lambda_{\mathfrak{p}})_{\mathrm{con}} \cong (\Lambda_{\mathrm{con}})_{\mathfrak{p}}$, as required.

(4) The proof is identical to (3). \square

4.2. The Contraction Theorem. In this subsection, so as to be able to work globally in future papers, we first relate the support of Λ_{con} to the locus L . We then control the support of Λ_{con} to deduce the deformation theory corollaries.

We need the following fact, which is well-known.

Lemma 4.5. *Suppose that $f: Y \rightarrow Z$ is a morphism of noetherian schemes which is not an isomorphism. Then $\mathbf{R}f_*: \mathrm{D}(\mathrm{Qcoh} Y) \rightarrow \mathrm{D}(\mathrm{Qcoh} Z)$ is not an equivalence.*

Proof. If $\mathbf{R}f_*$ is an equivalence then its inverse is necessarily given by its adjoint $\mathbf{L}f^*$. Since noetherian schemes are quasi-compact and quasi-separated, both unbounded derived categories are compactly generated triangulated categories (with compact objects the perfect complexes), and so the above equivalence restricts to an equivalence

$$\mathrm{per}(Y) \xleftarrow{\sim} \mathrm{per}(Z): \mathbf{L}f^*.$$

By Balmer [B, 9.7] it follows that f is an isomorphism, which is a contradiction. \square

Leading up to the next lemma, choose a closed point $\mathfrak{m} \in L$, and pick an affine open $\mathrm{Spec} R$ containing \mathfrak{m} . For any $\mathfrak{p} \in \mathrm{Spec} R$ we base change to obtain the following diagram.

$$\begin{array}{ccc} U_{\mathfrak{p}} & \xrightarrow{k} & U \\ \varphi_{\mathfrak{p}} \downarrow & & \downarrow f \\ \mathrm{Spec} R_{\mathfrak{p}} & \xrightarrow{j} & \mathrm{Spec} R \end{array}$$

Note that by flat base change the morphism $\varphi_{\mathfrak{p}}$ is still projective birational, and satisfies $\mathbf{R}\varphi_{\mathfrak{p}*} \mathcal{O}_{U_{\mathfrak{p}}} = \mathcal{O}_{R_{\mathfrak{p}}}$.

Lemma 4.6. *With the global setup of 3.1, and notation as above,*

$$\varphi_{\mathfrak{p}} \text{ is not an isomorphism} \iff \mathfrak{p} \in \mathrm{Supp}_R \Lambda_{\mathrm{con}}.$$

Proof. As in [KIWY, 4.6], it is easy to see that the diagram

$$\begin{array}{ccc} \mathrm{D}(\mathrm{Qcoh} U_{\mathfrak{p}}) & \xrightarrow[\sim]{\Psi := \mathbf{R}\mathrm{Hom}_{U_{\mathfrak{p}}}(\mathcal{O}_{U_{\mathfrak{p}}} \oplus k^* \mathcal{N}, -)} & \mathrm{D}(\mathrm{Mod} \Lambda_{\mathfrak{p}}) \\ \mathbf{R}\varphi_{\mathfrak{p}*} \downarrow & & \downarrow e(-) \\ \mathrm{D}(\mathrm{Qcoh} \mathrm{Spec} R_{\mathfrak{p}}) & \xlongequal{\quad\quad\quad} & \mathrm{D}(\mathrm{Mod} R_{\mathfrak{p}}) \end{array} \quad (4.I)$$

commutes, where the top functor is an equivalence by 4.4, and by abuse of notation e also denotes the idempotent in $\Lambda_{\mathfrak{p}}$ corresponding to $\mathcal{O}_{U_{\mathfrak{p}}}$.

(\Rightarrow) To ease notation, we drop \mathfrak{p} and write φ for $\varphi_{\mathfrak{p}}$. Since $\mathbf{R}\varphi_*$ is not an equivalence by 4.5, we may find some $x \in D(\mathrm{Qcoh} U_{\mathfrak{p}})$ such that the counit

$$\mathbf{L}\varphi^* \mathbf{R}\varphi_*(x) \xrightarrow{\varepsilon_x} x$$

is not an isomorphism, and so the object $c := \mathrm{Cone}(\varepsilon_x)$ is non-zero. Now we argue that $\mathbf{R}\varphi_*(c) = 0$, which is equivalent to the morphism $\mathbf{R}\varphi_*(\varepsilon_x)$ being an isomorphism. But

$$\mathbf{R}\varphi_*(x) \xrightarrow{\eta_{\mathbf{R}\varphi_*(x)}} \mathbf{R}\varphi_* \mathbf{L}\varphi^* \mathbf{R}\varphi_*(x) \xrightarrow{\mathbf{R}\varphi_*(\varepsilon_x)} \mathbf{R}\varphi_*(x)$$

gives the identity map by a triangular identity, and $\eta_{\mathbf{R}\varphi_*(x)}$ is an isomorphism since it is well known that $\eta: \mathrm{Id} \rightarrow \mathbf{R}\varphi_* \mathbf{L}\varphi^*$ is a functorial isomorphism by the projection formula. Thus indeed c is a non-zero object such that $\mathbf{R}\varphi_*(c) = 0$.

Now across the top equivalence in (4.1), since $c \neq 0$ it follows that $\Psi(c) \neq 0$ and so $H^j(\Psi(c)) \neq 0$ for some j . Further since the diagram (4.1) commutes, $e\Psi(c) = 0$, so since $e(-)$ is exact we deduce that $eH^i(\Psi(c)) = 0$ for all i . In particular there exists a non-zero $\Lambda_{\mathfrak{p}}$ -module $M := H^j(\Psi(c))$ such that $eM = 0$. It follows that M is a non-zero module for $(\Lambda_{\mathfrak{p}})_{\mathrm{con}}$, hence necessarily $(\Lambda_{\mathfrak{p}})_{\mathrm{con}} \neq 0$. But by 4.4 we have $(\Lambda_{\mathfrak{p}})_{\mathrm{con}} \cong (\Lambda_{\mathrm{con}})_{\mathfrak{p}}$, hence $\mathfrak{p} \in \mathrm{Supp}_R \Lambda_{\mathrm{con}}$.

(\Leftarrow) If $\mathfrak{p} \in \mathrm{Supp}_R \Lambda_{\mathrm{con}}$, by 4.4 $(\Lambda_{\mathfrak{p}})_{\mathrm{con}} \neq 0$. Hence the right hand functor $e(-)$ in (4.1) is not an equivalence, and so the left hand functor cannot be an equivalence either. By 4.5, it follows that $\varphi_{\mathfrak{p}}$ is not an isomorphism. \square

Theorem 4.7. *With the global setup of 3.1, choose a closed point $\mathfrak{m} \in L$, and pick an affine open $\mathrm{Spec} R$ containing \mathfrak{m} . Then*

- (1) $\mathrm{Supp}_R \Lambda_{\mathrm{con}} = L_R := L \cap \mathrm{Spec} R$.
- (2) $\mathrm{Supp}_{\mathfrak{R}} \Lambda_{\mathrm{con}} = \{\mathfrak{p} \in L_R \mid \mathfrak{p} \subseteq \mathfrak{m}\}$.

Proof. (1) It is clear that $L_R = \{\mathfrak{p} \in \mathrm{Spec} R \mid \varphi_{\mathfrak{p}} \text{ is not an isomorphism}\}$, and so the result is immediate from 4.6.

(2) Applying the above argument to the morphism $\mathfrak{U} \rightarrow \mathrm{Spec} \mathfrak{R}$, in an identical manner $\mathrm{Supp}_{\mathfrak{R}} \Lambda_{\mathrm{con}} = L_{\mathfrak{R}} := L \cap \mathrm{Spec} \mathfrak{R}$. It is clear that $L_{\mathfrak{R}} = \{\mathfrak{p} \in L_R \mid \mathfrak{p} \subseteq \mathfrak{m}\}$. \square

The following is an immediate corollary.

Corollary 4.8 (Contraction Theorem). *In the Zariski local setup $f: U \rightarrow \mathrm{Spec} R$ of 3.2, suppose further that $\dim U = 3$. Then*

$$f \text{ contracts curves without contracting a divisor} \iff \dim_{\mathbb{C}} \Lambda_{\mathrm{con}} < \infty.$$

Proof. In this setting f contracts curves without contracting a divisor if and only if L_R is a zero-dimensional scheme. The result follows from 4.7(1). \square

Remark 4.9. It follows from 4.8 (or indeed the morita equivalence in 3.7) that the condition $\dim_{\mathbb{C}} \Lambda_{\mathrm{con}} < \infty$ is independent of the choice of Λ , and thus the choice of tilting bundle of the form $\mathcal{O} \oplus \mathcal{N}$. Hence we may make any choice, and detect the contractibility by calculating the resulting $\dim_{\mathbb{C}} \Lambda_{\mathrm{con}}$. However, to get well-defined invariants that do not depend on choices, we pass to the formal fibre \mathfrak{R} and use the algebra Λ_{con} .

Corollary 4.10. *In the global setup $f: X \rightarrow X_{\mathrm{con}}$ of 3.1, suppose further that $\dim X = 3$, pick a closed point $\mathfrak{m} \in L$ and set $C := f^{-1}(\mathfrak{m})$. The following are equivalent.*

- (1) *There is a neighbourhood of \mathfrak{m} over which f does not contract a divisor.*
- (2) *The functor Def_X of simultaneous noncommutative deformations of the reduced fibre C^{red} is representable.*

Proof. By choosing an affine open $\mathrm{Spec} R$ containing \mathfrak{m} , this is identical to the proof of 4.8, appealing to 4.7(2) instead of 4.7(1), and using the prorepresentability of Def_X from 3.9. \square

5. DEFORMATIONS OF THE SCHEME-THEORETIC FIBRE

In the global setup $f: X \rightarrow X_{\text{con}}$ of 3.1, we choose a closed point $\mathfrak{m} \in L$ and in this section study commutative and noncommutative deformations of the scheme-theoretic fibre \mathcal{O}_C , where $C := f^{-1}(\mathfrak{m})$. We show that commutative and noncommutative deformations are prorepresented by the same object, and more remarkably that the prorepresenting object can be obtained from the same A as A_{con} can. This allows us to relate deformations of the reduced and scheme-theoretic fibres, in a way that otherwise would not be possible.

5.1. Prorepresentability. With the Zariski local setup in 3.2, taking the dual bundle in (3.A) induces a derived equivalence

$$\mathrm{D}^b(\mathrm{coh} U) \xrightarrow[\sim]{\mathbf{R}\mathrm{Hom}_U(\mathcal{V}^*, -)} \mathrm{D}^b(\mathrm{mod} \mathrm{End}_U(\mathcal{V}^*)). \quad (5.A)$$

Note that $\mathrm{End}_U(\mathcal{V}^*) = \Lambda^{\mathrm{op}}$. Also, by [V04, 3.5.7], under the above equivalence (5.A) the sheaf \mathcal{O}_C corresponds to a simple Λ^{op} -module, which we denote T'_0 . This fact is the reason we pass to \mathcal{V}^* , since it will allow us to easily apply 2.9 in the proof of 5.3 below.

Passing to the formal fibre $\mathfrak{U} \rightarrow \mathrm{Spec} \mathfrak{R}$, the dual of the previous bundle in 3.5 gives the following natural definition.

Definition 5.1. We write $\mathcal{M} := \bigoplus_{i=1}^n \mathcal{M}_i$ and define

$$B := \mathrm{End}_{\mathfrak{U}}(\mathcal{O}_{\mathfrak{U}} \oplus \mathcal{M}) = A^{\mathrm{op}},$$

which is the basic algebra morita equivalent to $\widehat{\Lambda}^{\mathrm{op}}$. From this, we define

$$B_{\mathrm{fib}} := \mathrm{End}_{\mathfrak{U}}(\mathcal{O}_{\mathfrak{U}} \oplus \mathcal{M})/[\mathcal{M}].$$

Under this dual setup, the following is obvious.

Lemma 5.2. $B_{\mathrm{fib}} \cong A_{\mathrm{fib}}^{\mathrm{op}} \cong A_{\mathrm{fib}}$, and in particular B_{fib} is commutative.

Proof. The first statement is clear since $B = A^{\mathrm{op}}$. The second statement is 3.6. \square

In what follows, we let S'_0 denote the simple B -module corresponding to T'_0 under the composition of the completion functor and the morita equivalence between $\widehat{\Lambda}^{\mathrm{op}}$ and B . Similarly to 3.8, for a given scheme-theoretic fibre C , below we use the following notation.

- (1) $\mathcal{D}ef_X^{\mathcal{O}_C}$ for the DG deformation functor associated to the injective resolution of $\mathcal{O}_C \in \mathrm{coh} X$.
- (2) $\mathcal{D}ef_U^{\mathcal{O}_C}$ for the DG deformation functor associated to the injective resolution of $\mathcal{O}_C \in \mathrm{coh} U$.
- (3) $\mathcal{D}ef_{\Lambda^{\mathrm{op}}}^{T'_0}$ for the DG deformation functor associated to the injective resolution of $T'_0 \in \mathrm{mod} \Lambda^{\mathrm{op}}$.
- (4) $\mathcal{D}ef_B^{S'_0}$ for the DG deformation functor associated to the injective resolution of $S'_0 \in \mathrm{mod} B$.

Theorem 5.3. In the global setup $f: X \rightarrow X_{\text{con}}$ of 3.1, pick a closed point $\mathfrak{m} \in L$ and set $C := f^{-1}(\mathfrak{m})$. Then

$$\mathcal{D}ef_X^{\mathcal{O}_C} \cong \mathrm{Hom}_{\mathbf{p}\mathrm{Art}_1}(A_{\mathrm{fib}}, -),$$

and further A_{fib} is commutative.

Proof. Exactly as in 3.9, we first claim that

$$\mathcal{D}ef_X^{\mathcal{O}_C} \cong \mathcal{D}ef_U^{\mathcal{O}_C} \cong \mathcal{D}ef_{\Lambda^{\mathrm{op}}}^{T'_0} \cong \mathcal{D}ef_B^{S'_0} \cong \mathrm{Hom}_{\mathbf{p}\mathrm{Art}_1}(B_{\mathrm{fib}}, -). \quad (5.B)$$

Under $i: U \hookrightarrow X$, since C is closed $\mathbf{R}i_*\mathcal{O}_C = i_*\mathcal{O}_C$, and so the first claimed isomorphism follows from 2.8. Under the derived equivalence (5.A) above, the sheaf \mathcal{O}_C corresponds to the module T'_0 , so the second claimed isomorphism follows from 2.9, exactly as in the proof of 3.9(2). The third claimed isomorphism follows from the fact that finite length Λ^{op} -modules supported at \mathfrak{m} are equivalent to finite length B -modules, and so the result follows from 2.8. For the last claimed isomorphism, since S'_0 is the vertex simple, as in [DW1, 3.1] it is clear that

$$\mathcal{D}ef_B^{S'_0} \cong \mathrm{Hom}_{\mathbf{Alg}_1}(B_{\mathrm{fib}}, -).$$

It remains to show that $B_{\text{fib}} \in \mathbf{pArt}_1$, but this holds since \mathfrak{R} , thus B , and thus B_{fib} , are complete with respect to their augmentation ideals. With (5.B) established, the remaining statement follows from 5.2. \square

5.2. Comparison of Deformations. One of the remarkable consequences of 3.9 and 5.3 is that there is a single A , of which various factors control different natural geometric deformation functors. Thus proving elementary facts for the ring A has strong deformation theory consequences; the following is one such example.

Proposition 5.4. *If $\dim_{\mathbb{C}} A_{\text{con}} < \infty$, then $\dim_{\mathbb{C}} A_{\text{fib}} < \infty$.*

Proof. Since $\mathcal{O}_{\mathcal{U}} \oplus \mathcal{M}$ is generated by global sections, by 4.3 $A \cong \text{End}_{\mathfrak{R}}(\mathfrak{R} \oplus f_*\mathcal{M})^{\text{op}}$. To ease notation we temporarily set $D = f_*\mathcal{M}$, so $A_{\text{con}}^{\text{op}} = \text{End}_{\mathfrak{R}}(\mathfrak{R} \oplus D)/[\mathfrak{R}]$ and $A_{\text{fib}}^{\text{op}} = \text{End}_{\mathfrak{R}}(\mathfrak{R} \oplus D)/[D]$. Since taking opposite rings does not affect dimension, in what follows we can ignore the ops.

To establish the result, we prove the contrapositive. If $\dim_{\mathbb{C}} A_{\text{fib}} = \infty$ then there exists a non-maximal prime ideal $\mathfrak{p} \in \text{Spec } \mathfrak{R}$ such that $(A_{\text{fib}})_{\mathfrak{p}} \neq 0$. As in 4.4 we have $(A_{\text{fib}})_{\mathfrak{p}} \cong (A_{\mathfrak{p}})_{\text{fib}}$, and thus $(A_{\mathfrak{p}})_{\text{fib}} = \text{End}_{\mathfrak{R}_{\mathfrak{p}}}(\mathfrak{R}_{\mathfrak{p}} \oplus D_{\mathfrak{p}})/[D_{\mathfrak{p}}] \neq 0$. Certainly this means that $D_{\mathfrak{p}}$ cannot be free. Now if $\text{Id}: D_{\mathfrak{p}} \rightarrow D_{\mathfrak{p}}$ factors through $\text{add } \mathfrak{R}_{\mathfrak{p}}$ then $D_{\mathfrak{p}}$ is projective. But since $\mathfrak{R}_{\mathfrak{p}}$ is local, $D_{\mathfrak{p}}$ would then be free, which is a contradiction. Thus $\text{Id}: D_{\mathfrak{p}} \rightarrow D_{\mathfrak{p}}$ does not factor through $\text{add } \mathfrak{R}_{\mathfrak{p}}$, so

$$(A_{\text{con}})_{\mathfrak{p}} \cong (A_{\mathfrak{p}})_{\text{con}} = \text{End}_{\mathfrak{R}_{\mathfrak{p}}}(\mathfrak{R}_{\mathfrak{p}} \oplus D_{\mathfrak{p}})/[\mathfrak{R}_{\mathfrak{p}}] \neq 0.$$

This implies that $\mathfrak{p} \in \text{Supp}_{\mathfrak{R}} A_{\text{con}}$ and so $\dim_{\mathbb{C}} A_{\text{con}} = \infty$. \square

Corollary 5.5. *In the global setup $f: X \rightarrow X_{\text{con}}$ of 3.1, pick a closed point $\mathfrak{m} \in L$ and set $C := f^{-1}(\mathfrak{m})$. Write $C^{\text{red}} = \bigcup_{i=1}^n C_i$, then*

- (1) *If Def_X is representable, so is $\text{Def}_X^{\mathcal{O}_C}$.*
- (2) *Suppose $\dim X = 3$. If $\text{Def}_X^{\mathcal{O}_C}$ is not representable, then f contracts a divisor over a neighbourhood of \mathfrak{m} .*

Proof. (1) This is immediate from 5.4, since A_{fib} prorepresents $\text{Def}_X^{\mathcal{O}_C}$ by 5.3, and A_{con} prorepresents Def_X by 3.9.

(2) Follows from (1) and 4.10. \square

The converse to 5.5(2) is however false; we show this in §6 below. The following summarises the main results in this paper in the case of 3-folds.

Summary 5.6. *In the global setup $f: X \rightarrow X_{\text{con}}$ of 3.1, suppose further that $\dim X = 3$, pick a closed point $\mathfrak{m} \in L$, set $C := f^{-1}(\mathfrak{m})$ and write $C^{\text{red}} = \bigcup_{i=1}^n C_i$.*

- (1) *Both the noncommutative deformation functor $\text{Def}_X^{\mathcal{O}_C}$ and commutative deformation functor $c\text{Def}_X^{\mathcal{O}_C}$ are prorepresented by A_{fib} .*
- (2) *The following are equivalent.*
 - (a) *The functors $c\text{Def}_X^{\mathcal{O}_C}$ and $\text{Def}_X^{\mathcal{O}_C}$ are representable.*
 - (b) $\dim_{\mathbb{C}} A_{\text{fib}} < \infty$.
- (3) *The following are equivalent.*
 - (a) *The functor Def_X of simultaneous noncommutative deformations of the reduced fibre C^{red} is representable.*
 - (b) $\dim_{\mathbb{C}} A_{\text{con}} < \infty$.
 - (c) *There is a neighbourhood of \mathfrak{m} over which f does not contract a divisor.*
- (4) *The statements in (3) imply the statements in (2), but the statements in (2) do not imply the statements in (3) in general.*

Proof. Part (1) is 5.3, and part (2) is tautological. Part (3) is 3.9 and 4.10, and part (4) is shown by the counterexample in 6.3 below. \square

6. EXAMPLES


In this section, we first illustrate some A_{con} that can arise in the setting of 4.10 for specific cA_n singularities. We then show that the converse to 5.5 is false, and also that 4.10 fails if we replace noncommutative deformations by commutative ones.

6.1. First Examples. Consider the cA_n singularities $\mathfrak{R} := \mathbb{C}[[u, v, x, y]]/(uv - f_1 \dots f_n)$ for some $f_i \in \mathfrak{m} := (x, y) \subset \mathbb{C}[[x, y]]$. The algebra A in 3.5, and thus the algebras A_{con} and A_{fib} , can be obtained using the calculation in [IW2, 5.29]. Here we make this explicit in two examples.

Example 6.1 (A 2-curve flop). Consider the case $f_1 = x$, $f_2 = y$ and $f_3 = x + y$. In this example there are six crepant resolutions of $\text{Spec } \mathfrak{R}$, and each has two curves above the origin. For one such choice, by [IW1, 5.2]

$$A = \text{End}_{\mathfrak{R}}(\mathfrak{R} \oplus (u, x) \oplus (u, xy)),$$

which by [IW2, 5.29] can be presented as the completion of the quiver with relations



given by the superpotential


$$W = \frac{1}{2}c_1a_1c_1a_1 + \frac{1}{2}c_2a_2c_2a_2 + \frac{1}{2}c_3a_3c_3a_3 + c_1c_2a_2a_1 - c_1a_1a_3c_3 - c_2c_3a_3a_2.$$

From this presentation, factoring by the appropriate idempotents it is immediate that $A_{\text{fib}} \cong \mathbb{C}$, and further

$$A_{\text{con}} \cong \begin{array}{c} \bullet \\ \xleftarrow{c_2} \\ 1 \end{array} \begin{array}{c} \xrightarrow{a_2} \\ \bullet \\ 2 \end{array} \quad \begin{array}{l} a_2c_2a_2 = 0 \\ c_2a_2c_2 = 0. \end{array}$$

Since A_{con} is finite dimensional, by 5.6(3) the contraction only contracts curves to a point.


Example 6.2 (A divisorial contraction). Consider the case $f_1 = f_2 = x$ and $f_3 = y$. In this case there are three crepant resolutions, and each has two curves above the origin. For one such choice, obtained by blowing up the ideal $u = v = x = 0$, the resolution is sketched as follows,




where above the origin there are two curves, and every other fibre over the y -axis contains only one curve. For this resolution

$$A = \text{End}_{\mathfrak{R}}(\mathfrak{R} \oplus (u, x) \oplus (u, xy))$$

which by [IW2, 5.29] can be presented as the completion of the quiver with relations



From this, we see immediately that $A_{\text{fib}} \cong \mathbb{C}[[y]]$, and A_{con} is the completion of the quiver




with no relations. Thus in this example both A_{fib} and A_{con} are infinite dimensional, which by 5.6 confirms that the contraction morphism contracts a divisor to a curve.

We remark that the above example, 6.2, also appears in [R83, 2.4] and [Z, 4.13].

6.2. Failure of Commutative Deformations. Here we give two more complicated examples. The first shows that the converse to 5.5(2) is false, and the second shows that 4.10 fails if we replace noncommutative deformations by commutative ones. In both examples, there is only one curve in the fibre above the closed point \mathfrak{m} .

Example 6.3 ($\text{Def}^{\mathcal{O}_C}$ does not detect divisors). Consider the group $G := A_4$ acting on its three-dimensional irreducible representation, and set $\mathfrak{R} := \mathbb{C}[[x, y, z]]^G$. It is well-known that in the crepant resolutions of $\text{Spec } \mathfrak{R}$, the fibre above the origin of $\text{Spec } \mathfrak{R}$ is one-dimensional: see [GNS, §2.4] and [NS].

For the crepant resolution given by $h: G\text{-Hilb} \rightarrow \text{Spec } \mathfrak{R}$, there are three curves above the origin meeting transversally in a Type A configuration. Further, in this case the tilting bundle from $G\text{-Hilb}$ has endomorphism ring isomorphic to the completion of the following McKay quiver with relations (see e.g. [LS, p13–14] [GLR, 5.2])



where ρ is a cube root of unity. By [W, 2.15], since there are two loops on the middle vertex we see that the middle curve is a $(-3, 1)$ -curve, and since there are no loops on the outer vertices, the outer curves are $(-1, -1)$ -curves. Also, by inspection

$$\text{End}_{\mathfrak{R}}(M_2)/[\mathfrak{R} \oplus M_1 \oplus M_3] \cong \frac{\mathbb{C}\langle\langle u, v \rangle\rangle}{\text{cl}(u^2, v^2)}, \quad (6.A)$$

where $\text{cl}(u^2, v^2)$ denotes the closure of the two-sided ideal (u^2, v^2) . Evidently, the above factor is infinite dimensional. Since there is a surjective map

$$\text{End}_{\mathfrak{R}}(M_2)/[\mathfrak{R}] \rightarrow \text{End}_{\mathfrak{R}}(M_2)/[\mathfrak{R} \oplus M_1 \oplus M_3]$$

it follows that $\text{End}_{\mathfrak{R}}(M_2)/[\mathfrak{R}]$ must also be infinite dimensional.

Now, contracting both the outer $(-1, -1)$ -curves in $G\text{-Hilb}$ we obtain a scheme \mathfrak{U} and a factorization

$$\begin{array}{ccc} G\text{-Hilb} & \xrightarrow{h} & \text{Spec } \mathfrak{R} \\ & \searrow & \uparrow f \\ & \mathfrak{U} & \end{array}$$

The example we consider is $f: \mathfrak{U} \rightarrow \text{Spec } \mathfrak{R}$. By construction, there is only one curve above the origin. As in [KIWY, 4.6]

$$\text{D}^b(\text{coh } \mathfrak{U}) \cong \text{D}^b(\text{mod } \text{End}_{\mathfrak{R}}(\mathfrak{R} \oplus M_2)).$$

Set $A := \text{End}_{\mathfrak{R}}(\mathfrak{R} \oplus M_2)$, then the quiver for A is obtained from the above McKay quiver by composing two-cycles that pass through M_1 and M_3 . In particular, in the quiver for A there is no loop at the vertex corresponding to \mathfrak{R} , so $A_{\text{fib}} = \mathbb{C}$. In particular, by 5.3, Def^{oc} is representable.

On the other hand $A_{\text{con}} = \text{End}_{\mathfrak{R}}(M_2)/[\mathfrak{R}]$, and we have already observed that this is infinite dimensional, so by 4.8 f contracts a divisor to a curve. It is also possible to observe this divisorial contraction using the explicit calculations of open covers in [NS].

Example 6.4 ($c\text{Def}^{\mathcal{J}}$ does not detect divisors). Consider again $\mathfrak{R} := \mathbb{C}[[x, y, z]]^G$ where G is the alternating group above in 6.3. Now, contracting instead the middle curve in $G\text{-Hilb}$ (instead of the outer curves we contracted above) gives a factorization

$$\begin{array}{ccc} G\text{-Hilb} & \xrightarrow{h} & \text{Spec } \mathfrak{R} \\ & \searrow g & \nearrow \\ & W & \end{array}$$

The original middle curve contracts to a closed point \mathfrak{m} in W , so picking an affine open $\text{Spec } T$ in W containing \mathfrak{m} , and passing to the formal fibre of g over this point, we obtain a morphism

$$g: \mathfrak{W} \rightarrow \text{Spec } \mathfrak{T}.$$

To this contraction, we associate the contraction algebra A_{con} using the procedure in 3.5. By [W, 3.5(2)], it follows from the uniqueness of prorepresenting object that

$$A_{\text{con}} \cong \text{End}_{\mathfrak{R}}(M_2)/[\mathfrak{R} \oplus M_1 \oplus M_3],$$

where we have recycled notation from 6.3. Hence by (6.A) we see that

$$A_{\text{con}} \cong \frac{\mathbb{C}\langle\langle u, v \rangle\rangle}{\text{cl}(u^2, v^2)}.$$

We have already observed that this is infinite dimensional, so g contracts a divisor by 4.8. Alternatively, we can see that g contracts a divisor by using the explicit open cover as in [NS, p40].

On the other hand, by general theory (see e.g. [DW1, 3.2]) $c\text{Def}^{\mathcal{J}}$ is prorepresented by the abelianization of A_{con} , which in this case is simply

$$A_{\text{con}}^{\text{ab}} \cong \frac{\mathbb{C}[[u, v]]}{(u^2, v^2)}.$$

By inspection, this is finite dimensional. This shows that the commutative deformation functor $c\text{Def}^{\mathcal{J}}$ is representable, even although a divisor is contracted to a curve.

REFERENCES

- [B] P. Balmer, *Presheaves of triangulated categories and reconstruction of schemes*, Math. Ann. **324** (2002), no. 3, 557–580.
- [BW] G. Brown and M. Wemyss, *Threefold flops via contraction algebra superpotentials*, in preparation.
- [DW1] W. Donovan and M. Wemyss, *Noncommutative deformations and flops*, arXiv:1309.0698, to appear Duke Math. J.
- [DW3] W. Donovan and M. Wemyss, *Twists and braids for general 3-fold flops*, arXiv:1504.05320.
- [ELO] A. Efimov, V. Lunts and D. Orlov, *Deformation theory of objects in homotopy and derived categories. I. General theory*. Adv. Math. **222** (2009), no. 2, 359–401.
- [E03] E. Eriksen, *An introduction to noncommutative deformations of modules*, Noncommutative algebra and geometry, 90–125, Lect. Notes Pure Appl. Math., **243**, Chapman and Hall/CRC, Boca Raton, FL, 2006.
- [E07] E. Eriksen, *Computing noncommutative deformations of presheaves and sheaves of modules*, Canad. J. Math. **62** (2010), no. 3, 520–542.
- [GM] W. Goldman and J. Millson, *The deformation theory of representations of fundamental groups of compact Kähler manifolds*, IHES Publ. Math. No. **67** (1988), 43–96.
- [GNS] Y. Gomi, I. Nakamura and K. Shinoda, *Hilbert schemes of G -orbits in dimension three*. Kodaira’s issue. Asian J. Math. **4** (2000), no. 1, 51–70.
- [GLR] B. Greene, C. I. Lazaroiu and M. Raugas, *D-branes on non-abelian threefold quotient singularities*, Nuclear Phys. B **553** (1999), no. 3, 711–749.
- [HT] Z. Hua and Y. Toda, *Contraction algebra and invariants of singularities*, arXiv:1601.04881.
- [IW1] O. Iyama and M. Wemyss, *Singular derived categories of \mathbb{Q} -factorial terminalizations and maximal modification algebras*, Adv. Math. **261** (2014), 85–121.

- [IW2] O. Iyama and M. Wemyss, *Reduction of triangulated categories and Maximal Modification Algebras for cA_n singularities*, arXiv:1304.5259, to appear Crelle.
- [KIWY] M. Kalck, O. Iyama, M. Wemyss and D. Yang, *Frobenius categories, Gorenstein algebras and rational surface singularities*, Compos. Math. **151** (2015), no. 3, 502–534.
- [K] Y. Kawamata, *On multi-pointed non-commutative deformations and Calabi-Yau threefolds*, arXiv:1512.06170.
- [L02] O. A. Laudal, *Noncommutative deformations of modules*. The Roos Festschrift volume, 2. Homology Homotopy Appl. **4** (2002), no. 2, part 2, 357–396.
- [LS] L. LeBruyn and S. Symens, *Partial desingularizations arising from non-commutative algebras*, arXiv:math/0507494.
- [M99] M. Manetti, *Deformation theory via differential graded Lie algebras*, Algebraic Geometry Seminars, 1998–1999 (Italian) (Pisa), 21–48, Scuola Norm. Sup., Pisa, 1999.
- [NS] A. Nolla de Celis and Y. Sekiya, *Flops and mutations for crepant resolutions of polyhedral singularities*, arXiv:1108.2352.
- [R83] M. Reid, *Minimal models of canonical 3-folds*, Algebraic varieties and analytic varieties (Tokyo, 1981), 131–180, Adv. Stud. Pure Math., 1, North-Holland, Amsterdam, 1983.
- [S08] E. Segal, *The A_∞ deformation theory of a point and the derived categories of local Calabi-Yaus*, J. Alg. **320** (2008), no. 8, 3232–3268.
- [T] Y. Toda, *Non-commutative width and Gopakumar–Vafa invariants*, Manuscripta Math. **148** (2015), no. 3–4, 521–533.
- [V04] M. Van den Bergh, *Three-dimensional flops and noncommutative rings*, Duke Math. J. **122** (2004), no. 3, 423–455.
- [W] M. Wemyss, *Aspects of the homological minimal model program*, arXiv:1411.7189.
- [Z] T. Zenger, *Contraction criteria for reducible rational curves with components of length one in smooth complex threefolds*, Pacific J. Math. **212** (2003), no. 2, 377–394.

WILL DONOVAN, KAVLI INSTITUTE FOR THE PHYSICS AND MATHEMATICS OF THE UNIVERSE (WPI),
 THE UNIVERSITY OF TOKYO INSTITUTES FOR ADVANCED STUDY, KASHIWA, CHIBA 277-8583, JAPAN.
E-mail address: will.donovan@ipmu.jp

MICHAEL WEMYSS, THE MAXWELL INSTITUTE, SCHOOL OF MATHEMATICS, JAMES CLERK MAXWELL
 BUILDING, THE KING'S BUILDINGS, PETER GUTHRIE TAIT ROAD, EDINBURGH, EH9 3FD, UK.
E-mail address: wemyss.m@googlemail.com